

## FAST TRACK

# Heterologous hybridization to a complementary DNA microarray reveals the effect of thermal acclimation in the endothermic bluefin tuna (*Thunnus orientalis*)

PEDRO C. CASTILHO,\* BRADLEY A. BUCKLEY,† GEORGE SOMERO\* and BARBARA A. BLOCK\*

\*Hopkins Marine Station, Stanford University, Pacific Grove, CA 93950, USA, †Department of Biology, Portland State University, Portland, OR 97201, USA

## Abstract

The temperature stress that pelagic fishes experience can induce physiological and behavioural changes that leave a signature in gene expression profiles. We used a functional genomics approach to identify genes that were up- or down-regulated following thermal stress in the Pacific bluefin tuna. Following the acclimation period, 113, 81 and 196 genes were found to be differentially expressed between the control (20 °C) and cold (15°) treatment groups, in ventricle, red muscle and white muscle, respectively. The genes whose expression levels were responsive to thermal acclimation varied according to muscle fibre type, perhaps reflecting the tissue-specific degrees of endothermy characteristic of this species.

*Keywords:* bluefin tuna, cDNA, cold, microarray, stress, transcriptome

*Received 16 October 2008; revision received 18 February 2009; accepted 27 February 2009*

## Introduction

Temperature is a major environmental factor that can govern the depth preferences and biogeographical distributions of pelagic fishes such as bluefin tunas. Their highly migratory behaviour puts bluefin tuna in contact with a wide range of ambient temperatures. Studies employing electronic tags describe bluefin tunas in highly diverse ocean environments, with depths of occurrence to over 1200 m and water temperatures ranging from 0.1 to 32 °C (Block *et al.* 2001; Kitagawa *et al.* 2007).

Physiological adaptations such as high cardiac output and regional endothermy make tunas an interesting model for studies in cardiac and locomotory muscle physiology and thermal stress responses. An elaborate vasculature system employing counter-current heat exchangers (*retia mirabilia*) in the viscera, brain and body musculature provides a means for the conservation of metabolic heat (Carey & Gibson 1983; Block & Carey 1985; Block 1991). The heat generated by muscle contraction and metabolism is thereby partially conserved, and the temperature of the internalized red muscle of bluefin

is maintained up to 21 °C above ambient water temperature (Carey & Lawson 1973; Marcinek *et al.* 2001). In contrast, the temperatures of heart and superficial white muscle remain at ambient water temperature. The different ranges of temperatures experienced by the endothermic and ectothermic regions of bluefin tunas raise a number of important questions about the thermal responses of the different types of muscle (red, white and cardiac) in these species.

Differences in thermal responses of these different muscle types are likely to be reflected in variations in patterns of gene expression, as indexed by mRNA populations (transcriptomes) of the cells. Global analysis of gene expression has proven to be a useful means to simultaneously assess the expression of tens of thousands of gene transcripts, providing a resolution and precision of phenotypic characterization not previously possible (Gracey 2007). By using this strategy, multiple genes and pathways can be examined as they work in concert. Using expressed sequence tag (EST) technology, several laboratories have generated complementary DNA (cDNA) microarrays of genes expressed in nonmodel species, including several species of fish (Gracey *et al.* 2001; Podrabsky & Somero 2004; Buckley *et al.* 2006). The use of microarrays for studying fish gene expression has increased in recent years, and arrays have emerged as an important tool for understanding developmental processes as well as basic physiology (Douglas 2006).

Correspondence: Pedro Castilho, University Federal Rural Pernambuco (UFRPE), DEPAq, Rua Dom Manoel de Medeiros, s/n Bairro de Dois Irmãos, CEP 52171-030, Recife, PE, Brazil. Fax: 55 81 33206512; E-mail: pccastilho@gmail.com

A growing number of studies are utilizing cDNA microarrays generated from one species to examine gene expression in other species – heterologous hybridization – to avoid the expense in time and money that is associated with designing novel microarrays for every new species of interest (Renn *et al.* 2004; Buckley 2007; Kassahn *et al.* 2007). Various factors have been considered that may affect the efficacy of this approach, including the phylogenetic distance between the species involved and the nature and length of DNA probes affixed to the microarray platform. Helpful in overcoming these issues is the choice of experimental design employed, comparing samples from a given species only against other samples from the same species.

In the current study, a cDNA microarray constructed from a Gobiid fish, the long jaw mudsucker *Gillichthys mirabilis* was used to generate gene expression profiles in the muscle tissues of a thermally acclimated Scombrid fish, the Pacific bluefin tuna (*Thunnus orientalis*). The *G. mirabilis* cDNA microarray was constructed from expressed sequence tags from liver, muscle and brain tissues from individuals that had been exposed to either heat or hypoxic stress, as well as from numerous control individuals (for complete descriptions of the cDNA microarray construction, see Gracey 2008). This platform therefore contains much of the stress-responsive genome of this species and has previously been employed in studies measuring the impact of environmental temperature on the *G. mirabilis* transcriptome (Buckley *et al.* 2006). Here, we utilized this genomic tool to characterize the transcriptomic response to thermal acclimation in three types of muscle tissues from bluefin tuna acclimated to either 15 °C (hereafter 'cold acclimated') or to 20 °C (hereafter 'control') to better understand the genomic underpinnings of the thermal physiology of these regionally endothermic fishes.

## Materials and methods

### *Experimental animal collection and thermal acclimation conditions*

Specimens were captured on barbless circle hooks off San Diego, California, in waters of 20–21 °C sea surface temperature. Following capture, the tuna were maintained in live wells on board the fishing vessel and transported over land by truck to the Tuna Research and Conservation Center (TRCC) at Stanford University's Hopkins Marine Station in Pacific Grove, California. At the TRCC, fish were held in 109 m<sup>3</sup> circular tanks containing seawater at 20 °C for 3 weeks. Four individuals were then transferred to 15 °C and held for 90 days while a second group of four individuals remained at 20 °C. The tunas were fed three times per week. Mean body size was 74.3 cm fork length (range 70–78 cm) and 8.30 kg (range 7.1–9.4 kg). Ventricle, white muscle and red muscle tissue (~500 mg) from *n* = 4 fishes from each group were sampled and stored at –80 °C until mRNA extraction was performed.

### *Total RNA and poly A-RNA extraction for hybridization*

Tuna tissues were homogenized with a Teflon pestle and total RNA was extracted according to a standard Trizol protocol (Invitrogen). RNA samples were quality checked by measuring A260/A280 ratios and confirmed by agarose gel electrophoresis. Detailed poly A RNA extraction procedures are available in Supplementary Material, Supporting information.

### *Microarray preparation*

The cDNA microarray prepared for the current study was constructed by arraying polymerase chain reaction (PCR)-amplified cDNA clones reverse transcribed from total RNA isolated from liver, skeletal muscle, and brain of the temperate goby *Gillichthys mirabilis*. The microarray consists of 9206 cloned ESTs. The *G. mirabilis* cDNA microarray construction is described in Gracey (2008).

### *Microarray hybridization*

Samples of 3 µg of mRNA (poly A RNA) from each tissue sampled were reverse transcribed to cDNA, using oligo(dT15) and pdN6 random hexamer primers and amino-allyl UTP. Incubation to 65 °C for 30 min in 0.2 mol/L NaOH and 0.1 mol/L EDTA was performed to remove the RNA template. Single-stranded cDNA was labelled with either Cy3 or Cy5 cyanine monoreactive dye (Amersham) prepared in 100 µL of dimethyl sulfoxide. The cDNA samples were incubated with 5 µL of either Cy3 or Cy5 for 1 h at room temperature and then cleaned through PCR purification columns (QIAGEN) and eluted in 10 µL of Sigma water. Samples were brought to a final volume of 40 µL in 25 mmol/L Hepes, 0.75 mg/mL tRNA (Sigma), 3× SSC and 0.2% SDS. Samples were incubated to 95 °C for 1 min, allowed to cool to room temperature for 5 min, and then applied to microarray slides. All the hybridizations were conducted overnight at 65 °C in hybridization chambers. After the hybridization, the slides were washed by immersion in 0.6× SSC and 0.03% SDS for 2 min to remove unbound dye and then dipped vigorously in 0.6× SSC wash solution. The slides were dried by centrifugation and scanned on an AXON GenePix 400B microarray scanner (Axon Instruments, Molecular Devices). Each individual sample was hybridized once with each dye (dye swap) having always one sample from the control- and one from the cold-acclimation group. The hybridizations performed are listed in Table 1.

### *Microarray data analysis*

The fluorescence ratio of Cy3 to Cy5 was extracted from the 24 arrays used in this experiment using GenePix Pro 4.0 software. This ratio reflects the relative intensity from control to cold-acclimated individuals, and vice versa, on dye swap

**Table 1** Array hybridization design used to compare cold stressed and control fishes

| Tissue       | Array  | Dye (fish ID)   | Treatment               |
|--------------|--------|-----------------|-------------------------|
| Ventricle    | (1252) | Cy3(1) × Cy5(2) | control × cold stressed |
|              | (1243) | Cy5(1) × Cy3(2) | control × cold stressed |
|              | (1250) | Cy3(3) × Cy5(4) | control × cold stressed |
|              | (1246) | Cy5(3) × Cy3(4) | control × cold stressed |
|              | (1232) | Cy3(5) × Cy5(6) | control × cold stressed |
|              | (1235) | Cy5(5) × Cy3(6) | control × cold stressed |
|              | (1283) | Cy3(7) × Cy5(8) | control × cold stressed |
|              | (1267) | Cy5(7) × Cy3(8) | control × cold stressed |
| Red Muscle   | (1227) | Cy3(1) × Cy5(2) | control × cold stressed |
|              | (1245) | Cy5(1) × Cy3(2) | control × cold stressed |
|              | (1231) | Cy3(3) × Cy5(4) | control × cold stressed |
|              | (1247) | Cy5(3) × Cy3(4) | control × cold stressed |
|              | (1233) | Cy3(5) × Cy5(6) | control × cold stressed |
|              | (1239) | Cy5(5) × Cy3(6) | control × cold stressed |
|              | (1255) | Cy3(7) × Cy5(8) | control × cold stressed |
|              | (1284) | Cy5(7) × Cy3(8) | control × cold stressed |
| White muscle | (1228) | Cy3(1) × Cy5(2) | control × cold stressed |
|              | (1242) | Cy5(1) × Cy3(2) | control × cold stressed |
|              | (1251) | Cy3(3) × Cy5(4) | control × cold stressed |
|              | (1248) | Cy5(3) × Cy3(4) | control × cold stressed |
|              | (1234) | Cy3(5) × Cy5(6) | control × cold stressed |
|              | (1240) | Cy5(5) × Cy3(6) | control × cold stressed |
|              | (1254) | Cy3(7) × Cy5(8) | control × cold stressed |
|              | (1285) | Cy5(7) × Cy3(8) | control × cold stressed |

$n = 4$  individuals were used for both control and cold stressed groups.

arrays. The data were visually inspected, and spots with low-quality areas of the array were flagged and excluded from further analysis. Spatial and intensity-based trends in the data were removed by Lowess normalization (GeneSpring, Agilent). For each spot on the array, the average value of the four control fish (value = control channel Cy5/reference channel Cy3) and their standard deviation were calculated. To test the cold-acclimated treatment, the experimental value (experimental channel Cy5/reference channel Cy3) for each spot on the array was tested if it fell outside two standard deviations of the mean for the control.

A *t*-test statistical analysis was used to identify genes for which the expression pattern showed a significant effect of the cold treatment. Different individuals ( $n = 4$ ) in each group were treated as replicates. For each spot on the array, the average value for all replicates and their standard deviation were calculated. Outliers were discarded and a new hybridization process was performed for at least two slides (regular and dye swap) for each fish tissue used in each treatment.

### Quantitative real-time PCR analysis: verification of microarray results

To confirm the expression levels generated from the microarray analysis, the expression patterns of a candidate gene was analysed using quantitative real-time PCR (RT-PCR) in a 96-well format (Bio-Rad). The Q-PCR assays were performed using iQ SYBR Green Supermix in accordance with manufacturers procedure (Bio-Rad) on an iQCyler platform. The cDNAs were synthesized from 1 µg of ventricle, red muscle or white muscle poly A RNA in a reaction of 25 µL using iScript cDNA Synthesis Kit (Bio-Rad) according to manufacturer's protocol. Template cDNA (1 µg/µL) was diluted 1:10:100 and run in triplicate. Control reactions without cDNA template were performed to check if amplified products were a result of DNA contamination or due to primer-dimer effects. Degenerate bluefin tuna primers were selected to amplify the candidate genes  $\beta$ -actin (EU 30041 – F 5'-CAAGAGAGGTATCCTGACC-3'; R 5' GTA GAAGGTGTGATGGGAG-3') and Hsp70 (EU 300943 F 5' GACATGAAGCACTGGC 3'; R 5' AGGACCATGGAGGAG 3'). The relative quantification  $2^{-\Delta\Delta CT}$  method described in Livak & Schmittgen (2001) was the method used to measure changes in the expression level of the target *Hsp70* gene and control housekeeping gene  $\beta$ -actin.  $\beta$ -actin was considered as a suitable housekeeping gene for normalization since its level of expression was independent of cold acclimation temperature [ $\Delta C_T \beta\text{-actin}$  (mean warm temp. – mean cold temp.) = 0]. One-way ANOVA analysis was performed to identify significant statistical differences in *Hsp70* relative gene expression levels between warm- and cold-acclimated bluefin tuna red muscle, ventricle and white muscle tissues. The Newman-Keuls test was applied a posteriori after the homogeneity of variances had been checked to the 95% confidence level.

### Results

Differences in gene expression between control- and cold-acclimated fish were examined using mRNA isolated from ventricle, white muscle and red muscle tissue. Table 2 lists those transcripts determined to be differentially expressed between the two groups that were successfully annotated by comparing their sequences to those in public databases. The number of differentially expressed genes varied among the three tissues. Seventy genes in ventricle, 36 genes in red muscle and 110 genes in white muscle were significantly up-regulated in the cold-acclimated group compared to the control group. Most genes that were up-regulated in ventricle in the cold acclimated group were involved with protein biosynthesis (14%) and transport (14%) (Table 2; Fig. 1a). Among the protein biosynthesis-related genes, the majority of the up-regulated genes encode the 60S ribosomal protein L-type. The up-regulated genes associated with transport encode a group of ferritin proteins, important for iron

**Table 2** Selected significantly differentially expressed transcript up- and down-regulated in red muscle, ventricle and white muscle of warm and cold acclimated BFT ( $P < 0.05$ )

| Up-regulated genes<br>Red muscle                           | Ventricle   | White muscle  |
|--|---|---|
| 60S acidic ribosomal protein P0                            | Glutamine/Q-rich-associated protein                         | (NC_003424) putative tata binding protein interacting protein |
| 60S ribosomal protein L19                                  | (NM_021245) myozenin 1; myozenin; calsarcin-2               | (NC_004331) hypothetical protein                              |
| 6-phosphofructokinase, liver type                          | 26S protease regulatory subunit 6B (TAT-binding protein-7)  | 26S proteasome non-ATPase regulatory subunit 1                |
| Alpha-parvin   | 26S proteasome non-ATPase regulatory subunit 1              | 40S ribosomal protein S11                                     |
| Amyloid protein-binding protein 1                          | 60S acidic ribosomal protein P0                             | 40S ribosomal protein S16                                     |
| ATP-binding cassette, subfamily B, member 7,               | 65 kDa Yes-associated protein                               | 40S ribosomal protein S19                                     |
| Cholesteryl ester transfer protein precursor               | Aldehyde oxidase  | 60S acidic ribosomal protein P1                               |
| Dystrobrevin binding protein 1 (Dysbindin)                 | ATP-dependent helicase DDX39                                | 60S ribosomal protein L10                                     |
| Glucosamine – fructose-6-phosphate aminotransferase        | BB282193  | 60S ribosomal protein L24                                     |
| <i>Homo sapiens</i> forkhead box P2                        | Calmodulin (CaM)  | 60S ribosomal protein L27                                     |
| Lipin 1  | CCAAT/enhancer binding protein zeta                         | 60S ribosomal protein L28                                     |
| L-lactate dehydrogenase A chain                            | Coagulation factor IX precursor                             | 60S ribosomal protein L34                                     |
| Next to BRCA1 gene 1 protein                               | Coronin-6 (Clipin E)  | 60S ribosomal protein L35a                                    |
| NK-tumour recognition protein                              | Cullin homologue 3 (CUL-3)                                  | 60S ribosomal protein L37                                     |
| Nucleolar transcription factor 1                           | Cyclin G1 (Cyclin G)  | 60S ribosomal protein L8                                      |
| Phosphatidylinositol transfer protein beta isoform         | Cysteine dioxygenase type I                                 | 65 kDa Yes-associated protein                                 |
| Phosphofurin acidic cluster sorting protein 1              | Cytochrome P450 24A1, mitochondrial precursor               | 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 4         |
| Phosphoribosyl pyrophosphate synthetase-associated protein | Desmoplakin (DP) 250/210 kDa paraneoplastic pemphigus       | Aldehyde oxidase  |
| Probable cation-transporting ATPase 13A3                   | Dihydroxyacetone phosphate acyltransferase                  | Alpha-actinin 1 (Alpha-actinin cytoskeletal isoform)          |
| Probable mannose-1-phosphate guanyltransferase             | Elongation factor 1-alpha (EF-1-alpha)                      | ATP-dependent Clp protease ATP-binding subunit ClpX-like      |
| Protein C20orf175  | Enigma homologue (Enigma-like PDZ and LIM domains protein)  | Beta-galactoside-binding lectin (14 kDa lectin)               |
| Putative adenosylhomocysteinase 2                          | Eukaryotic translation initiation factor 3 subunit 4        | Betaine – homocysteine S-methyltransferase                    |
| Putative polypeptide                                       | Eukaryotic translation initiation factor 3 subunit 7        | Cathepsin F precursor   |
| Retinol-binding protein IV, cellular                       | Ferritin, middle subunit (Ferritin M)                       | CC003931  |
| Ubiquitin  | Ferritin, middle subunit (Ferritin M)                       | PIR   T02995   T02995 unspecific monooxygenase-               |
| Ubiquitin carboxyl-terminal hydrolase isozyme L5           | Forkhead box protein P4 (Fork head-related protein-like A)  | Cell cycle control protein cwf5                               |
|  | Growth arrest and DNA-damage-inducible protein GADD45       | CG8947-PA ( <i>Drosophila melanogaster</i> )                  |
|  | GTP cyclohydrolase I feedback regulatory protein (GFRP)     | Chromosome 6 open reading frame 209 ( <i>Homo sapiens</i> )   |
|  | GTP:AMP phosphotransferase mitochondrial                    | Creatine kinase, M chain                                      |
|  | Guanine nucleotide-binding protein G(I)/G(S)/G(T) beta      | CTD small phosphatase-like protein (CTDSP-like)               |
|  | Heat shock cognate 71 kDa protein                           | Cytochrome P450 24A1, mitochondrial precursor                 |
|  | Histone H2B   | db55e12. × 1 Blackshear/Soares normalized <i>Xenopus</i>      |
|  |   | egg library <i>Xenopus</i>                                    |
|  | Hook homologue 2 (h-hook2)                                  | Density-regulated protein (DRP)                               |
|  | Hypothetical protein BC016816 ( <i>Homo sapiens</i> )       | Dipeptidyl-peptidase III (DPP III)                            |
|  | KIAA1040 protein ( <i>Homo sapiens</i> )                    | Elongation factor 1-alpha (EF-1-alpha)                        |
|  | Mid1 interacting protein 1 (Gastrulation specific G12-like) | Elongation factor 1-alpha, somatic form (EF-1-alpha-S)        |
|  | Mitochondrial carrier triple repeat 1                       | Eukaryotic translation initiation factor 3 subunit 1          |
|  | Myosin regulatory light chain 2, smooth muscle minor        | Eukaryotic translation initiation factor 3 subunit 7          |
|  | Nucleolar transcription factor 1 (Upstream binding)         | Eukaryotic translation initiation factor 4 gamma 2            |
|  | Probable dolichyl pyrophosphate Glc1Man9GlcNAc2             | Ferritin, middle subunit (Ferritin M)                         |
|  | Probable RNA-dependent helicase p68                         | fk94c05.y1 Zebrafish Research Genetics C32 fin Danio          |
|  |   | rerio cDNA 5'   |

Table 2 Continued


| Up-regulated genes<br>Red muscle | Ventricle  | White muscle  |
|----------------------------------|--|---|
|                                  | Procollagen-lysine,2-oxoglutarate 5-dioxygenase 1          | Gelsolin precursor (Actin-depolymerizing factor) (ADF)              |
|                                  | Proteasome subunit beta type 1-                            | Heat shock 70 kDa protein (HSP70)                                   |
|                                  | Protein FAM49B   | Heat shock cognate 71 kDa protein (Hsc70.1)                         |
|                                  | Protein KIAA1173   | Heat shock protein HSP 90-alpha                                     |
|                                  | Ras-related protein Rab-6B                                 | Heme oxygenase  |
|                                  | Similar to transcription factor UBF ( <i>Danio rerio</i> ) | hypothetical protein MGC63493 ( <i>Danio rerio</i> )                |
|                                  | Tubulin beta chain (Beta tubulin)                          | insulin-like growth factor binding protein 1 ( <i>Danio rerio</i> ) |
|                                  | Tubulin beta chain   | Jumonji/ARID domain-containing protein 1D (SmcY protein)            |
|                                  | Tyrosine-protein phosphatase, nonreceptor type 6           | Kelch-like protein 13 (BTB and kelch domain containing              |
|                                  | Ubiquitin  | Kelch-like protein 21   |
|                                  |  | KIAA0261 ( <i>Homo sapiens</i> )                                    |
|                                  |  | Maleylacetoacetate isomerase  |
|                                  |  | Mid1 interacting protein 1 (Gastrulation specific G12-like          |
|                                  |  | NipSnap2 protein  |
|                                  |  | Organic cation transporter protein                                  |
|                                  |  | Ornithine decarboxylase antizyme, long isoform                      |
|                                  |  | Parvalbumin alpha (A1)  |
|                                  |  | peptidylglycine alpha-amidating monooxygenase COOH-terminal         |
|                                  |  | Probable cysteine desulphurase, mitochondrial precursor             |
|                                  |  | Probable general secretion pathway protein E                        |
|                                  |  | Probable mitochondrial import receptor subunit TOM40                |
|                                  |  | Proteasome subunit alpha type 3                                     |
|                                  |  | Proteasome-associated protein ECM29 homologue                       |
|                                  |  | Protein C14orf103 homologue   |
|                                  |  | Protein expressed in T-cells and eosinophils in atopic dermatitis   |
|                                  |  | Ras-like protein 3 (Roughened protein)                              |
|                                  |  | Ras-related protein Rab-10  |
|                                  |  | Rho-related GTP-binding protein RhoG                                |
|                                  |  | Sarcalumenin precursor  |
|                                  |  | Serine - pyruvate aminotransferase, mitochondrial precursor         |
|                                  |  | Suppressor of G2 allele of SKP1 homologue                           |
|                                  |  | TC178346  |
|                                  |  | T-complex protein 1, zeta subunit                                   |
|                                  |  | T-complex protein 1, zeta subunit                                   |
|                                  |  | Tob1 protein  |
|                                  |  | Translocon-associated protein gamma subunit (TRAP-gamma)            |
|                                  |  | Ubiquitin   |
|                                  |  | Ubiquitin carboxyl-terminal hydrolase isozyme L5                    |
|                                  |  | Ubiquitin-conjugating enzyme E2 variant 2                           |
|                                  |  | VANGL2  vang-like 2 (van gogh, <i>Drosophila</i> ),                 |
|                                  |  | mRNA (cDNA clone MGC:76845)   |
|                                  |  | X box binding protein-1 (XBP-1)                                     |
|                                  |  | zgc:92423 ( <i>Danio rerio</i> )                                    |

Table 2 Continued

| Down-regulated genes<br>Red muscle                                   | Ventricle   | White muscle   |
|--|---|--|
| 26S proteasome non-ATPase regulatory subunit 3                       | (NC_002758) hypothetical protein ( <i>Staphylococcus aureus</i> ssp.) | 72 kDa type IV collagenase precursor                       |
| Actin, alpha skeletal muscle (Alpha-actin 1)                         | 40S ribosomal protein S20   | Acidic leucine-rich nuclear phosphoprotein 32 family       |
| ADP-ribosylation factor GTPase-activating protein 3                  | Alkaline phytoceramidase  | Actin, alpha skeletal muscle (Alpha-actin 1)               |
| Alpha-1 catenin (102 kDa cadherin-associated protein)                | ATP synthase alpha chain, mitochondrial precursor                     | Similar to SP   P07222   NPM_XENLA Nucleophosmin           |
| Apolipoprotein A-IV precursor  | BETA3 protein   | Aldehyde dehydrogenase family 7 member A1                  |
| Betaine – homocysteine S-methyltransferase                           | BJ047360 NIBB Mochii normalized <i>Xenopus</i> neurula library        | Alpha enolase (2-phospho-D-glycerate)                      |
| Brain-type organic cation transporter                                | CC162912  | Annexin A6 (Annexin VI) (Lipocortin VI)                    |
| Cytochrome P450 2J1  | CCAAT/enhancer binding protein delta (C/EBP delta)                    | Apolipoprotein A-IV precursor (Apo-AIV)                    |
| <i>Danio rerio</i> lipoprotein lipase mRNA                           | Creatine kinase, M chain  | ATP synthase alpha chain, mitochondrial precursor          |
| <i>Danio rerio</i> runt-related transcription factor b (runxb), mRNA | F-box only protein 32 (Muscle atrophy F-box protein)                  | ATP synthase beta chain, mitochondrial precursor           |
| Desmoplakin (DP) 250/210 kDa paraneoplastic pemphigus                | fj04h03.y1 Zebrafish adult olfactory <i>Danio rerio</i> cDNA 5', mRNA | Chitotriosidase 1 precursor                                |
| Dihydroceramide delta(4-desaturase)                                  | FK506-binding protein 1B  | Chymotrypsin B   |
| Fructose-bisphosphate aldolase A                                     | Fructose-bisphosphate aldolase A                                      | Citrate synthase, mitochondrial precursor                  |
| Haloacid dehalogenase-like hydrolase domain containing               | Heme oxygenase  | Cullin homologue 5 (CUL-5) (Vasopressin-activated)         |
| Hypothetical protein MGC33302 ( <i>Homo sapiens</i> )                | Insulin-like growth factor binding protein 1 ( <i>Danio rerio</i> )   | DNA-directed RNA polymerase II largest subunit             |
| Inhibitor of nuclear factor kappa-B kinase alpha subunit             | Intermediate filament protein ON3                                     | Enigma homologue (Enigma-like PDZ and LIM domains protein) |
| Microsomal signal peptidase 18 kDa subunit                           | Lipoprotein lipase precursor  | Enolase  |
| N-acetylglucosamine-6-sulphatase precursor                           | Myosin heavy chain, fast skeletal muscle                              | FK506-binding protein 1B                                   |
| Orphan nuclear receptor TR2  | Protein C20orf149   | Fructose-1,6-bisphosphatase                                |
| Oxysterol-binding protein 1  | Protein C20orf45 (CGI-107)  | Gastrulation specific protein G12                          |
| Pecanex-like 3 ( <i>Mus musculus</i> )                               | Protein kinase C and casein kinase substrate in neurones              | Glucose-6-phosphate isomerase                              |
| Polypyrimidine tract-binding protein 1 (PTB)                         | Serotransferrin I precursor (Siderophilin I) (STF I)                  | GLutamate dehydrogenase                                    |
| Probable aminotransferase T01B11.2                                   | Suppressor of cytokine signalling 3 (SOCS-3)                          | Glyceraldehyde-3-phosphate dehydrogenase                   |
| Protein kinase C, eta type   | Thrombospondin-1 precursor  | Glycogen phosphorylase, muscle form                        |
| Proto-oncogene protein c-fos (Cellular oncogene fos)                 | Tob1 protein (Transducer of erbB-2 1)                                 | Heat-shock protein beta-1 (HspB1)                          |
| Receptor-interacting serine/threonine-protein kinase 2               | TPR repeat containing protein KIAA1043                                | Importin beta-3 (Karyopherin beta-3)                       |
| SH3 domain GRB2-like protein B2 (Endophilin B2)                      | Tropomodulin-3 (Ubiquitous tropomodulin) (U-Tmod)                     | Intrinsic factor precursor (IF) (INF)                      |
| Similar to hypothetical protein ( <i>Rattus norvegicus</i> )         |   | Keratin, type II cytoskeletal 8 (Cytokeratin 8)            |
| Similar to hypothetical protein FLJ37794 ( <i>Homo sapiens</i> )     |   | KIAA0930 protein ( <i>Homo sapiens</i> )                   |
| Solute carrier family 2, facilitated glucose transporter,            |   | L-lactate dehydrogenase A chain                            |
| Thrombospondin-1 precursor   |   | Myosin heavy chain, fast skeletal muscle                   |
| Troponin C, slow skeletal and cardiac muscles                        |   | Myosin light chain 3, skeletal muscle                      |
| Tyrosine-protein phosphatase, nonreceptor type 2                     |   | Nicotinamide riboside kinase 2                             |
|  |   | Nucleolysin TIA-1 (RNA-binding protein TIA-1)              |


Table 2 Continued

| Down-regulated genes<br>Red muscle | Ventricle | White muscle  |
|------------------------------------|-----------|---|
|                                    |           | Nucleoside diphosphate kinase NBR-B<br>Pepsin A precursor<br>Phosphatidylinositol 3,4,5-trisphosphate-dependent<br>Phosphoglycerate mutase 2<br>Phosphorylase b kinase beta regulatory chain<br>Plasma protease C1 inhibitor precursor<br>Probable RNA-dependent helicase p68<br>Protein C14orf103<br>Protein C22orf19 (NF2/meningioma region protein pK1.3)<br>Protein FAM3A precursor<br>Protein FAM49B<br>Protein kinase C and casein kinase substrate in neurones<br>Proto-oncogene protein c-fos (Cellular oncogene fos)<br>Rabphilin-3 A (Exophilin 1)<br>Ribulose biphosphate carboxylase large chain precursor<br>RNA-binding protein 5 (RNA binding motif protein 5)<br>Sarcoplasmic/endoplasmic reticulum calcium ATPase 3<br>Serotransferrin I precursor (Siderophilin I) (STF I)<br>T-cell activation protein phosphatase 2C ( <i>Homo sapiens</i> )<br>Thrombospondin-1 precursor<br>Triosephosphate isomerase<br>Tropomyosin 1 alpha chain (Alpha-tropomyosin)<br>Troponin C, skeletal muscle (TNC)<br>Troponin T, fast skeletal muscle isoforms<br>Tumor differentially expressed protein 2 (Tumor)<br>Tyrosine aminotransferase<br>Unknown (protein for MGC:66340); wu:fb73e11 ( <i>Danio rerio</i> ) |




**Fig. 1** Percentage of total up-regulated genes identified in the Gene Ontologies (Biological Process) of ventricle (a), red muscle (b) and white muscle (c).

transport and homeostasis. An increase in the expression of a 71-kDa heat shock protein and cytochrome P450 were also observed. In the red muscle, genes encoding carbohydrate metabolism (14%), lipid metabolism (10%) and protein biosynthesis (10%) were also generally up-regulated (Fig. 1b). L-lactate dehydrogenase-A (LDH-A) and lipin 1, which are involved in carbohydrate and lipid metabolism, respectively, were both more highly expressed in the cold-acclimated group. As in the ventricle, an increase in the expression of genes encoding 60S ribosomal proteins was observed. Genes encoding products involved in protein biosynthesis (17%), protein folding (5%) and transport (9%) were up-regulated in white muscle from the cold-acclimation group (Fig. 1c). Several ribosomal proteins were up-regulated, including 40S and 60S ribosomal proteins (Table 2). Also up-regulated were heat shock genes encoding *Hsp70*, *Hsp71*, *Hsp90*.



**Fig. 2** Percentage of total down-regulated genes identified in the Gene Ontologies (Biological Process) of ventricle (a), red muscle (b) and white muscle (c).

Similar to the up-regulated genes, the number of differentially down-regulated genes varied among the three tissues tested. Following the cold-acclimation, 43 genes in ventricle tissue, 45 genes in red muscle tissue and 86 genes in white muscle were significantly down-regulated compared to the control group (Table 2). The most striking constituency of transcripts down-regulated in ventricle were those involved in transport (17%), regulation of transcription (8%), muscle development (8%) and regulation of cell growth (8%) (Fig. 2a). Genes encoding myosin heavy chain, lipase precursor and protein kinase C were among those that showed lower levels of expression in ventricle of cold-acclimated fish compared to control. Transcripts down-regulated in red muscle of cold-acclimated individuals were those involved in transport (13%), protein amino acid phosphorylation (11%) and DNA binding (8%) (Fig. 2b). Protein kinase C and



**Fig. 3** Relative gene expression level of candidate gene *Hsp70* in cold-stressed *Thunnus orientalis* ( $n = 7$ ) determined by quantitative real-time PCR. Expression levels were normalized to the house-keeping gene  $\beta$ -actin. (\*) Significant values between means.

cytochrome P450 were down-regulated in red muscle. Genes involved in carbohydrate metabolism (20%), transport (14%) and muscle development (8%) were down-regulated in white muscle (Fig. 2c). L-lactate dehydrogenase A was most prominently down-regulated among the genes involved in carbohydrate metabolism. The stress protein *Hsp70* was up-regulated in the cold-acclimated group only in the white muscle. This result was confirmed by an independent method, quantitative real-time PCR. The relative expression of *Hsp70* (normalized to  $\beta$ -actin) was significantly up-regulated in the cold-acclimated group only in white muscle, paralleling the results obtained via microarray analysis (Fig. 3).

## Discussion

This study examined the effect of thermal acclimation on broad-scale patterns of gene expression in endothermic and ectothermic muscle tissues of juvenile Pacific bluefin tuna. Each type of muscle tissue exhibited a distinct effect of acclimation on global gene expression, perhaps reflecting tissue-specific differences in thermal physiology and plasticity of response to variable environmental temperatures.

As is the case for most fishes, there is a general lack of genome sequence information for bluefin tuna. Therefore, we used a heterologous hybridization analysis using cDNA microarray generated from another fish species. The primary technical challenge presented by heterologous hybridization is the problem of sequence divergence between the species for which the microarray was constructed and the species providing the cDNA against which it was hybridized (Buckley 2007). The hybridization strategy we employed followed that of an earlier study designed to demonstrate the efficacy of heterologous hybridization. Renn *et al.* (2004) showed consistent expression profiles using a cDNA microarray that was generated from an African cichlid fish, *Astatotilapia burtoni*, to study the transcriptomes of seven other fish species from closely related taxa to a very distantly related species.

The current study avoids the problem of phylogenetic distance between *G. mirabilis* and *T. orientalis* because, in every case, samples from only one species (*T. orientalis*) were competitively hybridized against one another. Thus, the effect of phylogenetic distance between the target and probe cDNAs on hybridization kinetics affected each sample equally. The results of the present study support the ability of heterologous hybridization to reveal biologically relevant patterns of gene expression across considerable taxonomic distance in fishes. The fact that many transcripts in the differentially acclimated bluefin tuna were of unknown function demonstrates that, for nonmodel organisms, considerable potential exists for new environmentally sensitive genes to be discovered. It should be noted that another caveat associated with heterologous hybridization is that in some cases multiple or alternate members of a gene family will bind to a given feature on the array and so it is possible that discerning specifically which member of that gene family was detected by cross-species hybridization can potentially be an issue. Nevertheless, the grouping of differentially expressed genes into overall Gene Ontology categories does provide valuable information on the functional classes represented by these genes.

### *Differential gene expression in cardiac muscle following thermal acclimation*

Unlike tropical tuna species, bluefin tuna expanded their niche by migrating to higher latitudes where they experience sustained exposures to low temperatures (Block *et al.* 1998; Block *et al.* 2001). During migrations, bluefin tuna can spend long periods in 15 °C and lower water temperatures (Kitagawa *et al.* 2007). Adaptations to cold include elevated levels of endothermy and enhanced cardiac power output. Blank *et al.* (2004) demonstrated experimentally that bluefin tuna hearts maintain function at 2 °C. In a study comparing various tuna species, Castilho *et al.* (2007) demonstrated that the activity of SERCA2, a key cardiac enzyme, is highest in bluefin tuna when measured at 5 °C. While some fish species such as rainbow trout increase heart mass in response to cold-acclimation (Vornanen *et al.* 2005), relative heart mass of bluefin tuna does not change in response to cold (data not shown). Apparently, the bluefin tuna heart is able to adjust to low temperatures without having to increase the quantity of tissue or levels of cardiac enzymes. In accordance with this, cDNA microarray analyses in the current study indicate that none of the cardiac hypertrophy markers were up-regulated during cold acclimation.

### *Differential gene expression in red muscle following thermal acclimation*

Tunas have a higher proportion of slow-twitch, relative to fast twitch, myotomal muscle than other teleosts (Graham

*et al.* 1983). In tunas, as in all other fishes, slow sustained swimming is powered by slow-twitch aerobic red muscle, while fast-twitch glycolytic white muscle is recruited as swimming speed increases (Altringham & Johnston 1990). Evidence for adaptation to a higher operating temperature in deeper muscle tissues is reflected in the ability of deep tuna muscle to function at 30 °C, developing almost 40% more power than at 25 °C (Altringham & Block 1997).

Bluefin tuna red muscle comprises only a small portion of the total body mass but has a high fat content, high capillarity, high mitochondrial protein concentration and high myoglobin content, which results in the dark red colour (Korsmeyer *et al.* 1996). The microarray analysis indicated that bluefin tuna aerobic red muscle tissue increases expression of genes involved in carbohydrate and lipid metabolism in response to cold acclimation, possibly a process similar to the fattening observed during bluefin tuna migrations to oceanic cold water masses (Nottestad *et al.* 1999; Mourete *et al.* 2002).

In the current study, deep red muscle showed the lowest number of thermally responsive genes, possibly due to the temperature stability of this tissue, which works relatively independent of the external ambient temperatures.

#### *Differential gene expression in white muscle following thermal acclimation*

The white muscle, which comprises 46–55% of a bluefin tuna's body weight, and is known to accumulate unusually large amounts of lactate during burst activity (Dickson 1996). Acidification is avoided by rapidly removing the lactate after burst swimming (Guppy *et al.* 1979; Bernal *et al.* 2001). In response to cold acclimation, genes related to mitochondrial function and carbohydrate metabolism were down-regulated in white muscle (Table 2). However, the most dramatic effect of thermal acclimation on white muscle transcripts involved genes related to protein biosynthesis. The overall up-regulation of genes coding for ribosomal proteins suggests that the protein synthesis machinery of tuna white muscle is enhanced in the cold. Temperature related down-regulation of gene transcripts of carbohydrate metabolism suggests that while capacity for glycolytic energy production from carbohydrates might be decreased, mitochondrial function is not compensated.

In summary, gene expression patterns following thermal acclimation in bluefin tuna varied according to muscle fibre type, perhaps reflecting the tissue-specific degrees of endothermy characteristic of this species. The baseline data collected here may allow future studies to assess the thermal histories of wild populations of bluefin tuna that are acclimatized to different ranges of temperatures.

#### **Acknowledgements**

This research was supported by awards from the National Ocean Services/NOAA. The authors are grateful to indispensable technical

support of Chuck Farwell, Matt Price, Alexander Norton, Ty Brandt and Luis Rodriguez of TRCC.

#### **References**

- Altringham JD, Block BA (1997) Why do tuna maintain elevated slow muscle temperatures? Power output of muscle isolated from endothermic and ectothermic fish. *Journal of Experimental Biology*, **200**, 2617–2627.
- Altringham JD, Johnston IA (1990) Modeling muscle power output in a swimming fish. *Journal of Experimental Biology*, **148**, 395–402.
- Bernal D, Dickson KA, Shadwick RE, Graham JB (2001) Review: analysis of the evolutionary convergence for high performance swimming in lamnid sharks and tunas. *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative Physiology*, **129**, 695–726.
- Blank JM, Morrissette JM, Landeira-Fernandez AM *et al.* (2004) *In situ* cardiac performance of Pacific bluefin tuna hearts in response to acute temperature change. *Journal of Experimental Biology*, **207**, 881–890.
- Block BA (1991) Evolutionary novelties — how fish have built a heater out of muscle. *American Zoology*, **31**, 726–742.
- Block BA, Carey FG (1985) Warm brain and eye temperatures in sharks. *Journal of Computer Physiology*, **156**, 229–236.
- Block BA, Dewar H, Blackwell SB *et al.* (2001) Migratory movements, depth preferences, and thermal biology of Atlantic bluefin tuna. *Science*, **293**, 1310–1314.
- Block BA, Dewar H, Farwell C, Prince ED (1998) A new satellite technology for tracking the movements of Atlantic bluefin tuna. *Proceedings of the National Academy of Sciences, USA*, **95**, 9384–9389.
- Buckley BA (2007) Comparative environmental genomics in non-model species: using heterologous hybridization to DNA-based microarrays. *Journal of Experimental Biology*, **210**, 1602–1606.
- Buckley BA, Gracey AY, Somero GN (2006) The cellular response to heat stress in the goby *Gillichthys mirabilis*: a cDNA microarray and protein-level analysis. *Journal of Experimental Biology*, **209**, 2660–2677.
- Carey FG, Gibson QH (1983) Heat and oxygen-exchange in the *rete mirabile* of the bluefin tuna, *Thunnus thynnus*. *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative Physiology*, **74**, 333–342.
- Carey FG, Lawson KD (1973) Temperature regulation in free-swimming bluefin tuna. *Comparative Biochemistry and Physiology*, **44**, 375–&.
- Castilho PC, Landeira-Fernandez AM, Morrissette J, Block BA (2007) Elevated Ca<sup>2+</sup> ATPase (SERCA2) activity in tuna hearts: comparative aspects of temperature dependence. *Comparative Biochemistry and Physiology. Part A, Molecular & Integrative Physiology*, **148**, 124–132.
- Dickson KA (1996) Locomotor muscle of high-performance fishes: what do comparisons of tunas with ectothermic sister taxa reveal? *Comparative Biochemistry and Physiology*, **113A**, 39–49.
- Douglas SE (2006) Microarray studies of gene expression in fish. *Omics*, **10**, 474–489.
- Gracey AY (2007) Interpreting physiological responses to environmental change through gene expression profiling. *Journal of Experimental Biology*, **210**, 1584–1592.
- Gracey AY (2008) The *Gillichthys mirabilis* Cooper array: a platform to investigate the molecular basis of phenotypic plasticity. *Journal of Fish Biology*, **72**, 2118–2132.

- Gracey AY, Troll JV, Somero GN (2001) Hypoxia-induced gene expression profiling in the euryoxic fish *Gillichthys mirabilis*. *Proceedings of the National Academy of Sciences, USA*, **98**, 1993–1998.
- Graham JB, Koehn FJ, Dickson KA (1983) Distribution and relative proportions of red muscle in scombrid fishes – consequences of body size and relationships to locomotion and endothermy. *Canadian Journal of Zoology*, **61**, 2087–2096.
- Guppy M, Hulbert WC, Hochachka PW (1979) Metabolic sources of heat and power in tuna muscles. II. Enzymes and metabolite profiles. *Journal of Experimental Biology*, **82**, 303–320.
- Kassahn KS, Caley MJ, Ward AC *et al.* (2007) Heterologous microarray experiments used to identify the early gene response to heat stress in a coral reef fish. *Molecular Ecology*, **16**, 1749–1763.
- Kitagawa T, Boustany AM, Farwell CJ, Williams TD, Castleton MR, Block BA (2007) Horizontal and vertical movements of juvenile bluefin tuna (*Thunnus orientalis*) in relation to seasons and oceanographic conditions in the eastern Pacific Ocean. *Fish Oceanography*, **16**, 409–421.
- Korsmeyer KE, Dewar H, Lai NC *et al.* (1996) The aerobic capacity of tunas: adaptation for multiple metabolic demands. *Comparative Biochemistry and Physiology*, **113A**, 17–24.
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta C_T}$  method. *Methods*, **25**, 402–408.
- Marcinek DJ, Bonaventura J, Wittenberg JB, Block BA (2001) Oxygen affinity and amino acid sequence of myoglobins from endothermic and ectothermic fish. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, **280**, R1123–R1133.
- Mourete G, Megina C, Diaz-Salvago E (2002) Lipids in female northern bluefin tuna (*Thunnus thynnus*) during sexual maturation. *Fish Physiology and Biochemistry*, **24**, 351–363.
- Nottestad L, Giske J, Holst JC, Huse G (1999) A length-based hypothesis for feeding migrations in pelagic fish. *Canadian Journal of Fisheries and Aquatic Sciences*, **56**, 26–34.
- Podrabsky JE, Somero GN (2004) Changes in gene expression associated with acclimation to constant temperatures and fluctuating daily temperatures in an annual killifish *Austrofundulus limnaeus*. *Journal of Experimental Biology*, **207**, 2237–2254.
- Renn SCP, Aubin-Horth N, Hofmann HA (2004) Biologically meaningful expression profiling across species using heterologous hybridization to a cDNA microarray. *BMC Genomics*, **5**, 42.
- Vornanen M, Hassinen M, Koskinen H, Krasnov A (2005) Steady-state effects of temperature acclimation on the transcriptome of the rainbow trout heart. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, **289**, R1177–R1184.

---

P. C. Castilho is a Post-Doctoral Fellow studying the effect of cold temperatures in tunas. B. A. Buckley is interested in broad-scale patterns of environmentally controlled gene expression, as measured via cDNA microarrays. G. Somero studies physiological and biochemical adaptations of marine animals and B. A. Block is interested on evolutionary, cellular and molecular physiology.

---

### Supporting information

Additional supporting information may be found in the online version of this article:

#### Supplementary Material Methods

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.